

The Effect of 10 Weeks Resistance Training on Cholesterol and Blood Triglyceride Levels of Patients with Fatty Liver Disease

¹Rohollah Valizadeh, ²Siroos Hosseini Askarabadi, ³Sedigheh Karampour, ⁴Mona Abdolhamid Tehrani

ABSTRACT

The present study aims to consider the effect of 10 weeks resistance trainings on cholesterol and blood triglyceride (TG) levels of patients with having fatty liver, aged 50 to 60 in National Iranian South Oil Company (NISOC). This research is practical and its plan has been done experimentally with pre-test and post-test on experimental and control groups. In this study, 20 samples from 100 patients who referred to sonography clinic in NISOC with distinction of fatty liver were selected randomly and divided into two groups of control (n = 10) and experimental (n = 10). Cholesterol and blood triglyceride were measured as pretest. Test of normality for TG was (p = 0/200) by Kolmogorov-Smirnov and (p = 0/070) for cholesterol by Shapiro-Wilk test. After 10 weeks resistance trainings, the analysis and resolution of data were done by computer and SPSS (16) software as well as the descriptive and statistical methods (t-test). Comparison between these two groups showed that 8 weeks resistance trainings with a ≤ 0.05 causes significant decrease in the amount of TG but did not any significant effect on cholesterol of fatty liver patients.

Keywords: Resistance training, Cholesterol, Triglyceride, Fatty liver disease.

How to cite this article: Valizadeh R, Askarabadi SH, Karampour S, Tehrani MA. The Effect of 10 Weeks Resistance Training on Cholesterol and Blood Triglyceride Levels of Patients with Fatty Liver Disease. *Euroasian J Hepato-Gastroenterol* 2014;4(1):64-65.

Source of support: Nil

Conflict of interest: None declared

INTRODUCTION

Fatty liver disease and their complications represent significant public health problem in the world.^{2,4} Patients with fatty liver diseases express increased levels of cholesterol and triglycerides (TG). These patients are advised to undertake physical exercise¹⁻³ and one way to monitor prognosis of these patients may be accomplished by checking two important markers in the blood, cholesterol and TG. However, the role of physical exercises in regulating cholesterol and TG has not been fully elucidated in different parts of the world including Iran. Fatty liver diseases of these subjects were diagnosed by ultrasonography and from assessment of blood parameters. In this study, a total of 20 patients with fatty liver diseases working at National Iranian South Oil Company (NISOC), Iran, were enrolled. The ages of the patients varied from 50 to 60 years. The patients were divided in two groups: (1) patients receiving

no exercise (control), and (2) the other group underwent exercise. The intensity of exercise was altered from 50 to 95% for three sessions in these patients as shown in Figure 1.

There was no significant difference in levels of cholesterol among control or experimental patients during enrollment. Also, exercise did not have any significant effect on the levels of cholesterol (Table 1).

However, the levels of TGs decreased significantly due to exercise (Table 2).

It is not clear why exercise had an effect on TGs but not on cholesterol levels. Further study with altered intensities of exercise for prolonged duration would be required as an extension of this study.

ACKNOWLEDGMENT

The authors acknowledge the support of friends and colleagues of Islamic Azad University, Omidyeh Branch,

¹Department of Physical Education, Islamic Azad University, Omidyeh Branch, Omidyeh, Iran

²Department of Physical Education, Islamic Azad University, Behbahan Branch, Behbahan, Iran

^{3,4}Sama Technical and Vocational Training College, Islamic Azad University, Omidyeh Branch, Omidyeh, Iran

Address reprint requests to: Rohollah Valizadeh, Lecturer, Department of Physical Education, Islamic Azad University, Omidyeh Branch, Omidyeh, Iran, Phone: +98-916-986-8067, Fax: +98-652-322-2533, e-mail: valizadeh8328@gmail.com, valizadeh@iauo.ac.ir






Fig. 1: Exercise protocol

Table 1: Comparison of cholesterol in control and experimental groups

Stage	Variable	Groups	Number	Mean	Standard deviation	Significant level
Before exercise	Cholesterol	Experimental	10	192.2	31.90768	0.948
		Control	10	193.8	29.07003	
After exercise	Cholesterol	Experimental	10	176.9	25.07965	0.167
		Control	10	194.3	28.87540	

Table 2: Comparison of triglyceride in control and experimental groups

Stage	Variable	Groups	Number	Mean	Standard deviation	Degree of freedom	Amount of T	Significant level
Before exercise	Triglyceride	Experimental	10	223.4	18.33758	18	-0.086	0.933
		Control	10	224.1	18.18699			
After exercise	Triglyceride	Experimental	10	194.4	9.75477	18	-5.813	0.001
		Control	10	228	15.45603			

that helped them to do this project. Here, they should say that this research was supported by Islamic Azad University, Omidiyeh Branch (IAUOB).

REFERENCES

- Expert Panel. Summary of the second report of the national cholesterol education program (NCEP) expert panel on detection, evaluation and treatment of high blood cholesterol in adults. *JAMA* 1993;269(23):3015-3023.
- Falck-Ytter Y, Younossi ZM, Marchesini G, McCullough AJ. Clinical features and natural history of nonalcoholic steatosis syndromes. *Semin Liver Dis* 2001;21(1):17-26.
- Kemmler W, Engelke K, von Stengel S, Wienneck J, Laurer D, Kalender WA. Long-term 4-year exercise has a positive effect on menopausal risk factors: the Erlanger fitness osteoporosis prevention study. *J Strength Cond Res* 2007;21(1):232-239.
- Teli MR, James OFW, Burt AD, Bennett MK, Day CP. The natural history of nonalcoholic fatty liver: a follow-up study. *Hepatology* 1995;22(6):1714-1719.